

## **Global Ocean Internal Wave Database**

Victor Klemas  
Graduate College of Marine Studies  
University of Delaware  
Newark, DE 19716  
phone: (302) 831-8256 fax: (302) 831-6838 email: [klemas@udel.edu](mailto:klemas@udel.edu)

Quanan Zheng  
Graduate College of Marine Studies  
University of Delaware  
Newark, DE 19716  
phone: (302) 831-4698 fax: (302) 831-6838 email: [zheng@triton.cms.udel.edu](mailto:zheng@triton.cms.udel.edu)

Xiao-Hai Yan  
Graduate College of Marine Studies  
University of Delaware  
Newark, DE 19716  
phone: (302) 831-3694 fax: (302) 831-6838 email: [xiaohai@triton.cms.udel.edu](mailto:xiaohai@triton.cms.udel.edu)

Award # N00014-97-1-0648  
[http://www.onr.navy.mil/sci\\_tech/ocean/onrpgahj.htm](http://www.onr.navy.mil/sci_tech/ocean/onrpgahj.htm)

### **LONG-TERM GOALS**

Our long-term goal is to develop a global database of ocean internal waves observed primarily from spacecraft. This database will be publicly accessible and can be used for various purposes including understanding the statistical properties of internal waves and upper ocean dynamics in any ocean area of interest.

### **OBJECTIVES**

Our objectives are to extract information on ocean internal waves from Space Shuttle photographs and satellite SAR, to construct a database containing the information for global oceans and to make the database publicly accessible via the Internet. We will demonstrate the use of the database by performing statistical analysis of internal wave features and dynamic analysis of their evolution under continental shelf boundary conditions.

### **APPROACH**

The global database of ocean internal waves has two major sections, one for Space Shuttle images and a separate section containing SAR imagery from ERS-1, 2, Radarsat, and other spacecraft. The images will be accompanied by interpretation maps and text describing oceanographic properties of the imaged features. The database includes a home page, offers a standard format and is accessible to Internet users.

## WORK COMPLETED

We searched NASA and our own Space Shuttle archives to extract nearly 200 cases of ocean internal waves observed in various parts of the globe. A separate section of the database was set up for SAR images of internal waves, provided by various investigators. The radar section includes 50 SAR images. Interpretation maps and text describing oceanographic properties of the imaged features have been prepared to accompany one third of the images. A demonstration study for using the database has been performed, including statistical analysis of ocean internal wave features and dynamic analysis of the evolution of ocean internal waves under continental shelf boundary conditions.

## RESULTS

Based on the data contained in the database, one can conclude that ocean internal waves are common and can be observed by spaceborne SAR and visible imagery in all ocean regions, including the Atlantic, Pacific, Indian Oceans and major seas. In the open ocean their wavelengths have a Gaussian distribution, while closer to shore the distribution is skewed (Rayleigh). Most of the internal waves were generated by the interaction of the semidiurnal tide with the edge of the continental shelf. Tests showed that all internal wave images and interpretations maps in the database can be downloaded and accessed via the Internet. Adding a SAR section has significantly expanded the database to a total of about 250 scenes. Examples of a Space Shuttle image and a SAR image are shown in Figures 1 and 2, respectively.

Thermocline shoaling and deepening cause changes in the phase speeds of internal waves as described by linear wave theories. On the other hand, the ocean area where internal waves have variable phase speeds may be treated as a dynamically inhomogeneous medium. In this case, theories of nonlinear dispersive-wave propagation in inhomogeneous media may stand. We used these theories to analyze the evolution of ocean internal solitary waves passing over a seamount in the Gulf of Aden. The results indicate that a surprisingly sharp recess of an internal solitary wave packet, imaged by the space shuttle *Discovery*, is a signature of spatial phase delay caused by thermocline shoaling over the seamount. Soliton fission due to thermocline shoaling was also observed in the imagery. The observed number of transmitted solitons over the seamount agrees with theoretical predictions. Relative soliton amplitudes measured from the imagery also agree qualitatively with predictions.

We derived a theoretical model of a radar image for a KdV type ocean internal soliton, and validated the model using ocean internal wave signals taken from ERS-1 SAR and RadarSat SAR images. The results indicate that the model perfectly simulates ocean internal soliton signatures with double sign variations of radar backscatter. On the basis of the model, we developed the curve fitting method and the peak-to-peak method for determining the internal soliton characteristic half widths, which then were used to calculate the internal soliton amplitudes. The test results indicate that ocean internal soliton amplitudes derived by the two methods agree with in situ data acquired on the Portuguese Continental Shelf and the South China Sea with a reasonable accuracy. The role that wind fields play in ocean radar remote sensing was also analyzed. Finally, the modulation ratio of ocean internal waves on radar images was quantitatively estimated.

## IMPACT/APPLICATIONS

When completed, this Internet-accessible database will represent the largest collection of internal wave imagery observed by spacecraft over most of the globe. The sample size will be large enough for

scientists to evaluate the general statistical properties of internal waves in various parts of the oceans. Furthermore it will be possible to test models and obtain detailed descriptions of internal waves at specific ocean sites.

## TRANSITIONS

A dozen scientists from several countries have already requested imagery from our internal wave database. Nine papers using the data from this database have been published (or submitted) in scientific journals and conferences.

## RELATED PROJECTS

We have been working closely with various investigators, including Drs. John Apel (Global Ocean Associates), Antony Liu (NASA/GSFC), Tim Donato (NRL), etc. who have provided us with SAR imagery and performed some of the internal wave analysis.

## PUBLICATIONS

Klemas, V., Q. Zheng, and X.-H. Yan, 2000. A Database of Ocean Internal Waves. Proc. 28<sup>th</sup> Int. Symp. On Remote Sensing of Environment, Capetown, South Africa, March 27-30, 2000.

Klemas, V., Q. Zheng, and X.-H. Yan, 2000. Space Shuttle Studies of Ocean Internal Waves. Proc. Sixth Int. Conf. on Remote Sensing for Marine and Coastal Environments, Charleston, SC, May 1-3, 2000.

Zheng, Q., X.-H. Yan, W. T. Liu, and V. Klemas, 2000. Estimating Ocean Surface Currents Using Space Shuttle Images of Oil Slicks. AGU 2000 Ocean Sciences Meeting, San Antonio, TX, January 24-26, 2000.

Zheng, Q., X.-H. Yan, W. T. Liu, and V. Klemas, 2000. Solitary Waves in the Atmosphere and Ocean Observed from Space, Overseas Chinese Ocean-Atmosphere Remote Sensing Workshop, Hong Kong, China, July 6-7, 2000.



Zheng, Q., X.-H. Yan, W. T. Liu, V. Klemas, and D. Sun, Space shuttle observations of open ocean oil slicks. *Rem. Sens. Environ.*, 76, 49-56, 2001.

Zheng, Q., V. Klemas, X.-H. Yan, and J. Pan, Nonlinear evolution of ocean internal solitons as propagating along inhomogeneous thermocline. *J. Geophys. Res.*, 106, 14,083-14,094, 2001.

Klemas et al. Ocean Internal Wave Observations Using Space Shuttle and Satellite Imagery, *Geocarto International*, 16(2), 51-55, 2001

Zheng, G., Y. Yuan, V. Klemas, and X.-H. Yan, Theoretical expression for an ocean internal soliton SAR image and determination of the soliton characteristic half width, *J. Geophys. Res.*, 2001, in press.

Zheng, G., V. Klemas, and X.-H. Yan, Advance in studies of ocean internal waves observed from space, *Recent Research Developments in Geophysics*, Research Signpost, 2001 in press.



***Figure 1 Space shuttle images of internal waves in the Sulu Sea  
a) STS049-071-040 b) STS049-071-041 c) Interpretation map***

Two photographs were sequentially taken on May 8, 1992. The grouped curves are images of ocean internal waves in the Sulu Sea. There are six solitons in the wave packet. The observed portion of the leading wave crest is 190 km. The wavelength is about 10 km.

